

# USEFUL OVERVIEW IN ADOLESCENT CONTRACEPTION

---

NANCY SOKKARY 2019




---

No conflicts of Interest or disclosures

Nexplanon Trainer





**Congenital Anomalies of the Uterus/Vagina**  
**Congenital Malformations of the Vulva**  
**Transgender Adolescent Medicine**  
**Differences in Sexual Development**

### **Common Gynecologic Conditions**

Adnexal Masses, Cysts and Tumors

Contraception

Hirsutism

Genital Trauma

Sexually Transmitted Infections

Pediatric vulvar issues

Breast Abnormalities

### **Puberty and Menstruation**

Disorders of Puberty (Delayed or Precocious)

First Pelvic Exam

Menstrual suppression

Dysmenorrhea/Endometriosis

Menstrual Irregularities

Polycystic Ovary Syndrome

Premature Ovarian Insufficiency

(PMS)/ (PMDD)

# CONTRACEPTION: Objectives

---

General Principals

Counseling



Modes of Contraception

- Long acting reversible contraception
- Oral
- Non-Oral combined
- Injectable
  - \*efficacy based on weight

# Teen Pregnancy Outcomes

- Pregnancy Complications
  - Insufficient prenatal care
  - Low birth weight\*
  - Preterm Delivery\*
  - Less likely to breastfeed
- Future for mother
  - Postpartum depression
  - Less likely to graduate high school and go to college
  - Intimate partner violence
  - 18% with experience second pregnancy before age 20
- Future for infant
  - Developmental delay
  - More likely to be incarcerated
  - Poor school performance
  - Teen pregnancy

# Teen Pregnancy



Years,



<https://www.cdc.gov/teenpregnancy/about/index.htm>

<https://www.healthypeople.gov/2020/data/Chart/4467?category=1&by=Total&fips=-1>

GEORGIA CAMPAIGN FOR  
ADOLESCENT POWER & POTENTIAL  
**Education • Prevention • Action**  
*For Adolescent Health*



**DPH** Georgia Department of Public Health

We Protect Lives

Search this

About DPH Programs I Want To... Contact DPH

Programs

► District and County Operations

Home » Programs » Health Protection » Chronic Disease Prevention Section » Teen Pregnancy Prevention

## Teen Pregnancy Prevention

77° Login | Subscribe

**The Telegraph**

FULL MENU LOCAL NEWS SPORTS OBITUARIES TODAY'S DEALS

ALL DIGITAL ONLY \$ FOR 4 W

**SAMSUNG** Get the Galaxy S7 active for \$0 down\* with AT&T Next®, exclusively from AT&T.

\*Tax due at sale. Rec's well-qual. credit, elig. installment agmt & svc. If svc cancelled, device balance due. Activ. upgrade & other fees, charges & restr's apply. See \$0 down details. Device screen images simulated.



Learn more

EDUCATION

AUGUST 12, 2016 5:44 PM

## Sex ed, teen pregnancy to get bigger focus in new Bibb curriculum

# LAW

<b>Code Sections</b>	<b>Rape:</b> <a href="#">O.C.G.A. 16-6-1</a> <b>Statutory Rape:</b> <a href="#">O.C.G.A. 16-6-3</a>
<b>Elements of Rape</b>	<b>Rape:</b> Any man who forcibly uses his penis to penetrate a female's vagina against her will. This law only applies to females. A husband can rape his wife in Georgia.
<b>Elements of Statutory Rape</b>	<b>Statutory Rape :</b> Sexual intercourse with any person under the age of 16 years who is not your spouse. <b>Romeo and Juliet Provision:</b> If the victim is 14-16 years old and the defendant is either 18 years old or no more than four years older than the victim, he or she will be guilty of a misdemeanor.
<b>Penalty for Rape</b>	<a href="#">Death penalty</a> , life in prison without parole, or minimum of 25 years in prison followed by lifetime probation.
<b>Penalty for Statutory Rape</b>	<b>Statutory Rape: Felony</b> if under 21 years old, 1-20 yrs in prison. If over 21 years old, 10-20 yrs in prison and mandatory sex offender registration. <b>Romeo and Juliet Penalty: Misdemeanor</b> up to 1 yr. in jail, probation, fines, possible community service, and a possible "stay away" order.

**1-800-GACHILD**



# Laws

MINORS MAY CONSENT TO:						
STATE	CONTRACEPTIVE SERVICES	STI SERVICES	PRENATAL CARE	ADOPTION	MEDICAL CARE FOR MINOR'S CHILD	ABORTION SERVICES
Alabama	All†	All*	All	All	All	Parental Consent
Alaska	All	All	All		All	Parental Notice
Arizona	All	All		All		Parental Consent
Arkansas	All	All*	All		All	Parental Consent
California	All	All	All	All		▼ (Parental Consent)
Colorado	All	All	All	All	All	Parental Notice
Connecticut	Some	All		Legal counsel	All	All
Delaware	All*	All*	All*	All	All	Parental Notice‡
Dist. of Columbia	All	All	All	All	All	All
Florida	Some	All	All		All	Parental Notice
Georgia	All	All*	All	All	All	Parental Notice

# Medical Eligibility Criteria for Contraception

**Key:**


**1** No restriction (method can be used)

**2** Advantages generally outweigh theoretical or proven risks

**3** Theoretical or proven risks usually outweigh the advantages


**4** Unacceptable health risk (method not to be used)

The image displays two side-by-side tables, likely representing different versions or sections of the Medical Eligibility Criteria for Contraception. Each table has a header row with columns labeled 1 through 5. The rows represent various contraceptive methods. The cells are color-coded according to the key: green for 'No restriction (method can be used)', light green for 'Advantages generally outweigh theoretical or proven risks', pink for 'Theoretical or proven risks usually outweigh the advantages', and red for 'Unacceptable health risk (method not to be used)'. The tables are partially obscured by a black border on the left and right sides.



# Medical Eligibility Criteria

# Quick Start



# Condoms

---

Use them

The rate of dual use among adolescents is 22.8%, and is lowest among LARC users



# Counseling: AAP Policy

---



Pediatricians should counsel about and ensure access to a broad range of contraceptive services ...describing the most effective methods first.

**Contraception for Adolescents**  
COMMITTEE ON ADOLESCENCE  
*Pediatrics* 2014;134:e1244; originally published online September 29, 2014;  
DOI: 10.1542/peds.2014-2299

# Counseling

---

## **ACOG COMMITTEE OPINION**

Number 710 • August 2017


### **Committee on Adolescent Health Care**

*This Committee Opinion was developed by the American College of Obstetricians and Gynecologists' Committee on Adolescent Health Care in collaboration with committee member Karen R. Gerancher, MD.*

## **Counseling Adolescents About Contraception**

## EFFECTIVENESS OF FAMILY PLANNING METHODS\*

\*The percentages indicate the number out of every 100 women who experienced an unintended pregnancy within the first year of typical use of each contraceptive method.



Other Methods of Contraception: (1) Lactational Amenorrhea Method (LAM): is a highly effective, temporary method of contraception; and (2) Emergency Contraception: emergency contraceptive pills or a copper IUD after unprotected intercourse substantially reduces risk of pregnancy. Adapted from World Health Organization (WHO) Department of Reproductive Health and Research, Johns Hopkins Bloomberg School of Public Health/Center for Communication Programs (CCP), Knowledge for health project. Family planning: a global handbook for providers (2011 update). Baltimore, MD: Geneva, Switzerland: CCP and WHO; 2011; and Trussell J. Contraceptive failure in the United States. Contraception 2011;83:397-404.





# Bedsider.org

Are you a provider? Visit Bedsider Providers »

Welcome! (Sign in or Create your account) **BEDSIDERINSIDER**

**BEDSIDER**

birth control methods

where to get it

reminders


features

questions

search



# Stayteen.org



**STAYTEEN** ORG

SEX ED ASK US ANYTHING VIDEOS GAMES & QUIZZES CONNECT SEARCH

**Find a HEALTH CENTER**  
Enter zip code

**Get Protected**  
Curious about condoms? Interested in IUDs? Then get smart about all the best birth control that's out there. Compare methods, learn about side effects, and figure out which one is right for you!

**FILTER BY TOPIC**  
☐ BIRTH CONTROL / CONTRACEPTION ☐ ADVICE ☐ SEX ☐ RELATIONSHIPS ☐ ABSTINENCE  
RESET GO

**ARTICLE**  
Birth Control: Know ALL Your Options  
SHARES: 5

**ARTICLE**  
4 Things Every LGBTQ Teen Needs to Know  
SHARES: 11

**ARTICLE**  
I Have Epilepsy. Am I Loveable? Can I Have Sex?  
SHARES: 2

# Contraception

Most common type of contraception used by adolescent=least effective

- Combined Oral Contraceptive
- Condom
- With drawl

## Contraceptive Effectiveness

Proportion of women who will become pregnant over one year of use, by method

Method	Perfect use	Typical use
Implant	0.05	0.05
Vasectomy (male sterilization)	0.10	0.15
Intrauterine device (IUD)		
Levonorgestrel-releasing	0.2	0.2
Copper-T	0.6	0.8
Tubal (female) sterilization	0.5	0.5
Injectable	0.2	6
Pill	0.3	9
Vaginal ring	0.3	9
Patch	0.3	9
Diaphragm	6	12
Sponge**	9/20	12/24
Male condom	2	18
Female condom	5	21
Withdrawal	4	22
Fertility awareness methods***	0.4–5	24
Spermicides	18	28
Emergency contraception	*	*
No method	85	85

Notes: u = unavailable. "Perfect use" denotes effectiveness among couples who use the method both consistently and correctly; "typical use" refers to effectiveness experienced among all couples who use the method (including inconsistent and incorrect use). \*The effectiveness of emergency contraception (EC) is not measured on a one-year basis like other methods. EC is

# Contraception

---

Long Acting Reversible Contraception=LARC

- Implantable method
- IUDs
  - Copper, Mirena, Skyla, Liletta, Kyleena



# Contraception


---

Continuation rate of LARC at 12 months=81%

Continuation rate of short acting methods=47%

Failure rate of LARC=0.05-.08%


Failure rate of short acting combined methods=1-9%



<http://www.choiceproject.wustl.edu/>

Diedrich JT, Klein DA, Peipert JF. Long-acting reversible contraception in adolescents: a systematic review and meta-analysis. *Am J Obstet Gynecol* 2017;216:364.e12.

# Duration


**Table 1.** Long-Acting Reversible Contraceptive Methods 

Brand Name	Medication and Device Type (Dose)	Initial Rate of Release (micrograms/day)	FDA-approved Duration of Use	Potential Efficacy Beyond FDA-approved Duration	Identifying Characteristics	Size of Device (Horizontal x Vertical, mm)	Inserter Tube Diameter (mm)	Percentage of Women Experiencing an Unintended Pregnancy in the First Year of Use (Typical Use)*
Kyleena	LNG-IUD (19.5 mg)	17.5	5 years	N/A	Blue strings; silver ring	28 x 30	3.8	0.20 <sup>†</sup>
Liletta	LNG-IUD (52 mg)	19.5	4 years	+1 year <sup>‡</sup>	Blue strings	32 x 32	4.4	0.20 <sup>†</sup>
Mirena	LNG-IUD (52 mg)	20	5 years	+2 years <sup>§,  </sup>	Gray strings	32 x 32	4.4	0.20 <sup>†</sup>
Skyla	LNG-IUD (13.5 mg)	14	3 years	N/A	Gray strings; silver ring	28 x 30	3.8	0.20 <sup>†</sup>
Paragard	Copper T380A IUD (380 mm <sup>2</sup> )	NA	10 years	+2 years <sup>§</sup>	White strings	32 x 36	4.01	0.80
Nexplanon/ Implanon	Etonogestrel single-rod contraceptive implant (68 mg)	60–70	3 years	+1–2 years <sup>¶</sup>	N/A	40 x 2	N/A	0.05

# Implantable method

---

- Effective for 3-5 years
- 68mg etnogesterel
- No SS difference in efficacy based on weight
- Decrease blood loss and dysmenorrhea
- No effect on bone mineral density
- Specific training required
- BREAKTHROUGH BLEEDING



# Breakthrough bleeding

---

- COUNSELING
- NSAIDs
  - High dose 5-7 days
- Combined oral contraceptive
  - 20-35mcg pill 1-3 months
- Estrogen
  - 2mg estradiol po x 7 days
- Doxycycline
  - 100mg BID x5 days
- Tranexamic Acid
  - 650mg BID x5da



# Intrauterine Device

---



## Safety in adolescent

- No increase risk of infertility
- May be inserted without difficulty
- Possible increased risk of expulsion
- Infection
  - Risk of PID 0-5%
  - Highest if active infection when IUD is placed



# Intrauterine Device

Levonorgestrel: 5-7 year

- Mirena
- Liletta
- Effective for treating HMB, dysmenorrhea, endometriosis, hyperplasia
- No difference based on weight

5 year

- Kyleena\*

3 year

- Skyla\*



# Intrauterine Device

---

## Copper

- 10-12 years
- Hormone free
- May increase pain/cramping
- May be used as emergency contraception



# Oral Contraception

---

## Combined

- 50mcg
- 30mcg
- Lo-dose
- Drospirenone containing
- Triphasic

Progesterone only\*\*\*

No difference in efficacy among obese patients (kind of)

# Combined COC

---

## **Sprintec/Orthocyclin**

- Monophasic: .3/.35 estrogen/varying types of progesterone

## **Necon, Ovral**

- Monophasic: 50mcg
- Refractory bleeding

# Low dose pill

---

## Loestren, Lo-LoEstren

- 10-25mcg of estrogen
- May have decrease side effects
- Efficacy is equivalent
- +/- increased BTB

# Drospirenone containing COC

---

Yasmin, Yaz, Gianvi, Ocella, Zarah

- PCOS
- Has anti-androgenic affect
- Small diuretic affect
- PMS/PMDD
- 24/4 cycle

FDA: association with increase r/o thrombo-embolism but data not strong enough to conclude causality

# Triphasic

Why?

- No difference in breakthrough bleeding
- No difference in discontinuation rate



# Breast Cancer Update

Prospective Cohort study of 1.8 million Danish women

- Followed for 11 years
- Relative risk among current or recent users;
  - 1.20 [1.14 to 1.26]
  - 1 extra breast cancer per 7960 women per year
  - May be higher with longer use
  - May be lower if used < 5 years

**Table 4.** Relative Risk of Breast Cancer among Women Using Various Types of Hormonal Contraception Who Were Followed until December 31, 2012.\*

Variable	No. of Person-Yr	No. of Breast-Cancer Events	Age-Adjusted Incidence Rate <i>no. of events/100,000 person-yr</i>	Adjusted Relative Risk (95% CI)†	Age-Adjusted Risk Difference (95% CI) <i>no. of events/100,000 person-yr</i>
Never used hormonal contraception	7,815,180	5955	55	1.00 (Reference)	Reference
Used hormonal contraception >6 mo previously	4,348,722	2883	58	1.08 (1.03 to 1.13)	3 (1 to 6)
<b>Current or recent use of combined hormonal contraception</b>					
Oral combined ethinyl estradiol, 50 µg					
Norethisterone	52,895	23	46	1.01 (0.67 to 1.52)	-9 (-30 to 12)
Levonorgestrel	73,125	54	64	1.21 (0.93 to 1.59)	9 (-9 to 27)
Oral combined ethinyl estradiol, 20 to 40 µg					
Norethisterone	153,603	39	67	1.09 (0.80 to 1.50)	12 (-12 to 35)
Levonorgestrel	638,936	380	72	1.33 (1.20 to 1.48)	17 (9 to 25)
Norgestimate	635,732	180	72	1.22 (1.20 to 1.48)	18 (5 to 30)
Desogestrel	1,453,690	368	64	1.12 (1.01 to 1.25)	9 (1 to 17)
Gestodene	2,633,355	705	69	1.20 (1.11 to 1.30)	14 (8 to 20)
Drospirenone	503,700	102	60	1.05 (0.86 to 1.28)	6 (-8 to 20)
Cyproterone	272,804	77	90	1.44 (1.15 to 1.81)	36 (11 to 60)
Estradiol valerate and dienogest	6,380	7	101	1.62 (0.77 to 3.41)	46 (-30 to 122)
Nonoral combined hormonal contraception					
Patch	10,842	2	60	0.85 (0.21 to 3.41)	5 (-1 to 11)
Vaginal ring	91,313	20	53	0.97 (0.62 to 1.50)	-2 (-32 to 28)
<b>Current or recent use of progestin-only products</b>					
Oral contraceptive					
Norethisterone	128,848	78	58	1.00 (0.80 to 1.25)	3 (-10 to 16)
Levonorgestrel	10,547	16	102	1.93 (1.18 to 3.16)	47 (-4 to 99)
Desogestrel	77,847	42	69	1.18 (0.87 to 1.60)	14 (-8 to 36)
Nonoral contraceptive					
Implant	42,217	9	46	0.93 (0.48 to 1.79)	-9 (-42 to 25)
Levonorgestrel-releasing intrauterine system	503,441	571	70	1.21 (1.11 to 1.33)	16 (9 to 22)
Depot medroxyprogesterone acetate	19,308	5	51	0.95 (0.40 to 2.29)	-4 (-49 to 42)

\* Recent use was defined as discontinuation of hormonal contraception within the previous 6 months.

† Relative risks were adjusted for age, calendar year, level of education, the polycystic ovary syndrome, endometriosis, parity, and family history of premenopausal breast or ovarian cancer.




# Non-contraceptive benefits of COC

---

- Dysmenorrhea
- Cycle control
- Reduces menstrual blood loss by 40-50%
- Treatment of Menstrual Migraines\*
- Improve hirsutism and acne
- may decrease risk of colorectal, endometrial and ovarian cancer

# Helping continuation



# Combination Contraception

---

## Contraceptive Ring: Nuvaring

- As or more effective than OCP
- No difference in efficacy based on weight\*
- Have patient place it in clinic
- Maybe associated with vaginal irritation, expulsion, discomfort during sex

## Contraceptive patch: ortho-evra/xulane

- As or more effective than COCs
- Maybe less effective in obese women\*
- Slightly increased r/o thromboembolism compared to COC
  - Data contradictory

\*Obese Women: Subset analysis

# Medroxyprogesterone acetate (Depo)

---

## Weight gain

- Over 36 months gained 5.1kg more than OC users
- Only contraception regularly associated with weight gain

## BMD

- Decrease in BMD
- Longer use worsens BMD and may increase r/o fracture
- Reversible in younger patients

Consider in patients with Sick Cell or Seizures

No difference in efficacy among obese women

# Emergency Contraception

---

Over the counter\*



## Emergency Contraception

---

Levonorgestrel 0.75 mg- take 1 tab and repeat in 12 hours OR  
Levonorgestrel 1.5- 1 tab in a single dose

- Up to 72 hours post intercourse\*
- Failure rate up to 3 %
- \$50

# Emergency Contraception

---

## Ella/ulipristal- 1 tab single dose

- Up to 120 hours post intercourse
- Failure rate 1.4%
- \$51




# Emergency contraception

---

## Copper IUD

- Up to 120 hours post intercourse
- Failure rate  $<1\%$





# Questions?

---

# Emergency Contraception

## COC

- 2-4 pills repeat in 12 hours
- Efficacy not well studied

Twenty-one brands of oral contraceptives that can be used for emergency contraception (EC) in the United States

Brand	Company	Pills per dose*	Ethinyl Estradiol per dose, microgram	Levonorgestrel per dose, mg•
Plan B <sup>Δ</sup>	Barr	1 white pill	0	0.75
Ovral	Wyeth-Ayerst	2 white pills	100	0.50
Ogestrel	Watson	2 white pills	100	0.50
Cryselle	Barr	4 white pills	120	0.60
Levora	Watson	4 white pills	120	0.60
Lo/Ovral	Wyeth-Ayerst	4 white pills	120	0.60
Low-Ogestrel	Watson	4 white pills	120	0.60
Levlen	Berlex	4 light orange pills	120	0.60
Nordette	Wyeth-Ayerst	4 light orange pills	120	0.60
Portia	Barr	4 pink pills	120	0.60
Seasonale	Barr	4 pink pills	120	0.60
Trivora	Watson	4 pink pills	120	0.50
Tri-Levlen	Berlex	4 yellow pills	120	0.50
Triphasil	Wyeth-Ayerst	4 yellow pills	120	0.50
Enpresse	Barr	4 orange pills	120	0.50
Alesse	Wyeth-Ayerst	5 pink pills	100	0.50
Lessina	Barr	5 pink pills	100	0.50
Levite	Berlex	5 pink pills	100	0.50
Lutera	Watson	5 white pills	100	0.50
Aviane	Barr	5 orange pills	100	0.50
Ovrette	Wyeth-Ayerst	20 yellow pills	0	0.75
Jolessa	Barr	4 pink pills	120	0.60
Lybrel	Wyeth-Ayerst	6 yellow pills	120	0.54
Quasense	Watson	4 white pills	120	0.60
Seasonique	Duramed	4 blue-green pills	120	0.60

\* The treatment schedule is one dose as soon as possible after unprotected intercourse, and another dose 12 hours later. However, recent research has found that both doses of Plan B or Ovrette can be taken at the same time.

• The progestin in Cryselle, Lo/Ovral, Low-Ogestrel, Ogestrel, Ovral, and Ovrette is norgestrel, which contains two isomers, only one of which (levonorgestrel) is bioactive; the amount of norgestrel in each tablet is twice the amount of levonorgestrel. Levonorgestrel regimens also can be formulated by substituting double the amount of norgestrel as is indicated for levonorgestrel.

Δ Plan B is the only dedicated product specifically marketed for emergency contraception in the United States. Preven, a combined emergency contraception pill, is no longer available for the US market.

Not-2-Late.com: the emergency contraception website. Princeton University Office of Population Research, Princeton (NJ). Reproduced with permission from: ACOG Practice bulletin #69: Emergency Contraception. Obstet Gynecol 2005; 106:1443. Copyright © 2005 Lippincott Williams & Wilkins.