




CYSTIC FIBROSIS NEWBORN SCREENING

IMPORTANT FINDINGS FOR CARE PROVIDERS

Cystic fibrosis (CF) newborn screening (NBS) has been performed throughout the United States since 2010. CF newborn screening has led to early diagnosis in most affected infants and improves health outcomes. A summary of evaluations on the first decade of universal NBS is below.

CF NEWBORN SCREENING LEADS TO EARLY DIAGNOSIS IN MANY AFFECTED INFANTS.

65% of infants with a positive NBS test had an age of first event* within the first 30 days, as recommended by guidelines, allowing for earlier diagnosis and intervention. [2, 4]




*surrogate measure for timeliness inclusive of sweat test, clinic visit, and/or hospitalization

1 OF EVERY 8


infants with CF born between 2010-2018 were evaluated **after two months**.

EARLIER EVALUATION FOR CF IS ASSOCIATED WITH BETTER EARLY LIFE NUTRITION.



Why is this important?
Better early life nutrition is associated with improved lung function and survival. [3, 4, 5, 8, 9, 11]


LUNG FUNCTION HAS IMPROVED SINCE NBS IMPLEMENTATION IN 2010.



DETECTION OF CFTR VARIANTS VARIES ACROSS RACE AND ETHNICITY

Since 2020, all U.S. Cystic Fibrosis NBS programs have used CFTR gene variant detection. Detection of **at least one** variant defines a positive test and requires follow-up. Some states detect more variants than others, but all miss more infants from non-white races and Hispanic ethnicity. [6, 7]

AVERAGE DETECTION OF 1 CFTR VARIANT ACROSS 9 COMMON VARIANT PANELS BY RACE AND ETHNICITY



“

DELAYED DIAGNOSIS AND FALSE NEGATIVE NBS CONTINUE TO IMPACT PEOPLE WITH CF.

11%

of people with CF are estimated to have experienced a delayed diagnosis (>180 days) or been diagnosed after a false-negative NBS. [11]



BLACK/AFRICAN AMERICAN & ASIAN POPULATIONS EXPERIENCE THE HIGHEST RATE OF FALSE-NEGATIVE NBS OR DELAYED DIAGNOSIS AMONG NON-WHITE POPULATIONS. [1]


”

REFERRAL FOR EVALUATION AND TREATMENT SHOULD NOT BE DELAYED IN INFANTS WITH A POSITIVE NBS TEST.

40% of infants had weight-for-age percentiles below the 10th percentile at first CF Center visit from 2010-2018. [4]



CF CENTER & NBS PROGRAM-REPORTED BARRIERS



1

TIMELY EVALUATION AND TREATMENT IMPROVES HEALTH OUTCOMES.

Newborns with a positive NBS should be referred for a sweat test as soon as possible and no later than 28 days. Sweat testing can be done if weight is at or greater than 2 kilograms [4.4 pounds]. [2]



2

INFANTS CAN HAVE CF EVEN WITH A NEGATIVE NBS. INFANTS WITH CLINICAL CONCERN SHOULD BE REFERRED FOR FURTHER EVALUATION.

False negative NBS and the detection of one - or no - variants is possible in all infants. Infants showing clinical signs, including bowel obstruction and failure to thrive, within the first month of life should be referred for evaluation, even with a negative NBS or detection of only one variant. [2]



3

CLEAR COMMUNICATION CAN REDUCE MISCONCEPTIONS AND IMPROVE UNDERSTANDING OF RISK AND OUTCOMES.

There are long held misconceptions that CF only affects infants of European ancestry. All infants can have CF. Ensuring that health providers and caregivers understand this and the benefit of earlier diagnosis is essential.



Ann & Robert H. Lurie
Children's Hospital of Chicago
Stanley Manne
Children's Research Institute

**CYSTIC FIBROSIS
FOUNDATION**
ADDING TOMORROWS

This project was supported by the Centers for Disease Control and Prevention of the U.S. Department of Health and Human Services (HHS) as part of a financial assistance award totaling \$400,000 with 100 percent funded by CDC/HHS. The contents are those of the author(s) and do not necessarily represent the official views of, nor an endorsement, by CDC/HHS, or the U.S. Government. Funding for the research cited was provided by Cystic Fibrosis Foundation (MCCOLL1600).

1.Cystic Fibrosis Foundation Patient Registry 2021 Annual Data Report Bethesda, Maryland ©2022 Cystic Fibrosis Foundation

2.Farrell, P. M., White, T. B., Ren, C. L., Hempstead, S. E., Accurso, F., Derichs, N., Howenstone, M., McColley, S. A., Rock, M., Rosenfeld, M., Sermet-Gaudelus, I., Southern, K. W., Marshall, B. C., & Sosnay, P. R. (2017). Diagnosis of cystic fibrosis: Consensus guidelines from the Cystic Fibrosis Foundation. The Journal of Pediatrics, 181. <https://doi.org/10.1016/j.jpeds.2016.09.064>

3.Kronstad, M. W., Butler, S. M., Wohl, M. E., Stoddard, M., Matousek, R., Wagoner, J. S., Johnson, C. A., & Morgan, W. J. (2003). Growth and nutritional indexes in early life predict pulmonary function in cystic fibrosis. The Journal of Pediatrics, 142(6), 624-630. <https://doi.org/10.1016/j.mp.2003.152>

4.Martiano, S. L., Elbert, A. A., Farrell, P. M., Ren, C. L., Santag, M. K., Wu, R., & McColley, S. A. (2020). Outcomes of infants born during the first 9 years of CF newborn screening in the United States: A retrospective cystic fibrosis foundation patient registry cohort study. Pediatric Pulmonology, 56(12), 3758-3767. <https://doi.org/10.1002/ppul.25658>

5.Martiano, S. L., Elbert, A. A., Farrell, P. M., Fisher, N., Ren, C. L., Santag, M. K., Wu, R., & McColley, S. A. (2020). P022 earlier contact with a cystic fibrosis centre is associated with better nutritional outcomes in infants with cystic fibrosis. Journal of Cystic Fibrosis, 19. <https://doi.org/10.1016/j.jcf.2020.03.059-3>

6.McColley, S. A., Martiano, S. L., Ren, C. L., Santag, M. K., Rychlik, K., Bolmert, L., Elbert, A. A., Wu, R., & Farrell, P. M. (2023). Disparities in first evaluation of infants with cystic fibrosis since implementation of newborn screening. Journal of Cystic Fibrosis, 22(1), 89-97. <https://doi.org/10.1016/j.jcf.2022.07.010>

7.McGarry, M. E., Ren, C. L., Wu, R., Farrell, P. M., & McColley, S. A. (2022). Detection of disease-causing CFTR variants in state newborn screening programs with nutritional and pulmonary outcomes. JAMA Pediatrics, 176(10), 990. <https://doi.org/10.1001/jamapediatrics.2022.2674>

8.Rosenfeld, M., Ostregg, J., Cromwell, E. A., Wagret, A., Szczepaniak, R., Fink, A., Schechter, M. S., Fano, A., Ren, C. L., Morgan, W., & Sanders, D. B. (2022). Real-world associations of US cystic fibrosis newborn screening programs with nutritional and pulmonary outcomes. JAMA Pediatrics, 176(10), 990. <https://doi.org/10.1001/jamapediatrics.2022.2674>

9.Sanders, D. B., Fink, A., Mayer-Hamblett, N., Schechter, M. S., Sawicki, G. S., Rosenfeld, M., Fume, P. A., & Morgan, W. J. (2015). Early life growth trajectories in cystic fibrosis are associated with pulmonary function at age 6 years. The Journal of Pediatrics, 167(5). <https://doi.org/10.1016/j.jpeds.2015.07.044>

10.Santag, M. K., Miller, J. I., McKasson, S., Gaviglio, A., Martiano, S. L., West, R., Vazquez, M., Ren, C. L., Farrell, P. M., McColley, S. A., & Keller-Guenther, Y. (2022). Newborn screening for cystic fibrosis: A qualitative study of successes and challenges from Universal Screening in the United States. International Journal of Neonatal Screening, 8(3), 38. <https://doi.org/10.3390/ijns803038>

11.Yen, E. H., Quinton, H., & Borowitz, D. (2019). Better nutritional status in early childhood is associated with improved clinical outcomes and survival in patients with Cystic Fibrosis. The Journal of Pediatrics, 162(3). <https://doi.org/10.1016/j.jpeds.2012.08.040>